

AD-A114 054

NEW YORK UNIV NY DEPT OF CHEMISTRY

F/G 20/8

CHARGE TRANSFER SPECTRA AND PHOTOELECTRON EMISSION BY SOLUTIONS--ETC(U)

APR 82 P DELAHAY

N00014-75-C-0397

UNCLASSIFIED

TR-19

NL

1 of 1  
42-A  
114054

END

DATE

FILED

05-82

DTIC

AD A114054

12

OFFICE OF NAVAL RESEARCH  
Contract N00014-75-C-0397  
Task No. NR 051-258  
TECHNICAL REPORT NO. 19 (New Series)

CHARGE TRANSFER SPECTRA AND PHOTOELECTRON  
EMISSION BY SOLUTIONS  
by  
Paul Delahay

Accepted for publication in  
Chemical Physics Letters

New York University  
Department of Chemistry  
New York, NY

April 1982

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release  
and sale; its distribution is unlimited

DTIC  
ELECTE  
APR 29 1982  
H

82 04 29 038

DTIC FILE COPY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 19 (New Series)	2. GOVT ACCESSION NO. AD-A114054	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CHARGE TRANSFER SPECTRA AND PHOTOELECTRON EMISSION BY SOLUTIONS		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Paul Delahay		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0397
9. PERFORMING ORGANIZATION NAME AND ADDRESS New York University New York, NY 10003		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 051-258
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217		12. REPORT DATE April 1982
		13. NUMBER OF PAGES 22
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES  Accepted for publication in Chemical Physics Letters.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Acceptor, electron Charge transfer spectrum Donor, electron Photoelectron emission Reorganization free energy		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) → Charge transfer absorption spectra are correlated to the energetics of photoelectron emission by donor and/or acceptor solutions. Donor- acceptor pairs include: 10 inorganic anions and solvent (water); five cations ( $V^{2+}$ , $Co^{2+}$ ), and solvent; solvent and atoms (H, Cl, Br, I), OH radical, $Fe^{3+}$ , $Ce^{4+}$ ; anion-cation pairs exhibiting charge transfer in the 7 to 10 eV range. ←		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

## CHARGE TRANSFER SPECTRA AND PHOTOELECTRON EMISSION BY SOLUTIONS

PAUL DELAHAY

Department of Chemistry, New York University, New York, New York 10003, U.S.A.

Charge transfer absorption spectra are correlated to the energetics of photoelectron emission by donor and/or acceptor solutions. Donor-acceptor pairs include: 10 inorganic anions and solvent (water); five cations ( $V^{2+}$  to  $Co^{2+}$ ) and solvent; solvent and atoms (H, Cl, Br, I), OH radical,  $Fe^{3+}$ ,  $Ce^{4+}$ ; anion-cation pairs exhibiting charge transfer in the 7 to 10 eV range.

Optical donor-acceptor charge transfer (CT) will be correlated to photoelectron emission by solutions of the donor. The interpretation involves the calculation of the free energies for the CT and emission processes and the consideration of the reorganization free energies [1,2] on the basis of the theories of electron transfer [2] of Marcus [3] and Hush [4].

### 1. Basic relationships

The free energy  $\Delta G^{CT}$  for donor-acceptor CT will be obtained first. The donor and acceptor are represented by  $D^-$  and A, respectively, without consideration of the actual ionic charges (to simplify notations). Consider the following sequence: The separated reactants  $D^-$  and A in solution are brought within a distance at which CT occurs; the CT process is governed by the Franck-Condon principle, and CT therefore is followed by reorganization of the nuclear configurations of the products of CT and the solvent; finally, the products D and  $A^-$  are separated.

The preceding sequence is equivalent to the net reaction  $D^- + A = D + A^-$  involving the change of free energy  $\Delta G(DA^-)$ . Hence, one has [1,2]

$$\Delta G^{CT} = \Delta G(DA^-) - R^{CT}, \quad (1)$$

where  $R^{CT}$  ( $< 0$ ) is the reorganization free energy for the change of nuclear

configuration following CT. The work terms for bringing together reactants and separating products (weak bonding) are omitted in (1) as they are generally negligible ( $< 0.1$  eV) in the experiments considered here.

The contributions of the donor and acceptor in (1) will be separated. One sets,

$$\Delta G(DA^-) = \Delta G(D^-) + \Delta G(A), \quad (2)$$

where  $\Delta G(D^-)$  and  $\Delta G(A)$  are the changes of free energies for the reactions  $D^- + H^+ = D + 1/2H_2(g)$  (oxidation of  $D^-$ ) and  $A + 1/2H_2(g) = A^- + H^+$  (reduction of A) in solution, respectively.

The reorganization free energy  $R^{CT}$  in (1) will be interpreted on the basis of the theories of electron transfer in [3,4]. The quantity  $R^{CT}$  is equated to the sum of inner- and outer-sphere contributions (denoted by "in" and "out"). The outer-sphere contribution is derived from the macroscopic model of non-equilibrium polarization of a continuous medium [3]. Thus,


$$R_{out}^{CT} = -(\epsilon_{op}^{-1} - \epsilon_s^{-1})e^2(1/2a_D + 1/2a_A - 1/r_{DA}), \quad (3)$$

where  $\epsilon_{op}$  and  $\epsilon_s$  are the optical and static dielectric constants of water, respectively;  $a_D$  and  $a_A$  are the radii of the assumed spherical boundaries between inner and outer regions of the donor and acceptor, respectively;  $r_{DA}$  is the distance between the centers of the donor and acceptor prevailing in CT. The fields of the donor and acceptor in (3) are assumed to have spherical symmetry without mutual influence [2]. This approximation will suffice here. Equation (3) is rewritten in the form,

$$R_{out}^{CT} = R_{out}(D_*) + R_{out}(A_*) + R(DA^-), \quad (4)$$

where the asterisks denote the non-equilibrium nuclear configurations prior to reorganization.

One obtains from (1), (2) and (4) by noting that reorganization free energies are the sum of inner- and outer-sphere contributions,



$$\Delta G^{CT} = \Delta G(D^-) - R(D_*) + \Delta G(A) - R(A_*^-) - R(DA^-). \quad (5)$$

This equation is combined with the relationship [1,2,5] for the free energy  $\Delta G^m(D^-)$  for photoelectron emission by aqueous solutions of  $D^-$ ,

$$\Delta G^m(D^-) = \Delta G_H + \Delta G(D^-) - R(D_*), \quad (6)$$

where  $\Delta G_H = 4.50$  eV on the assumption of a negligible surface potential ( $\pm 0.1$  V) at the water vapor-solution interface. Thus,

$$\Delta G^m(D^-) - \Delta G_H = \Delta G^{CT} - \Delta G(A) + R \quad (7)$$

with

$$\begin{aligned} R &= R(A_*^-) + R(DA^-) \\ &= R_{in}(A_*^-) + R_{out}(A_*^-) + R(DA^-). \end{aligned} \quad (8)$$

Equation (7) correlates the energetics of CT and emission by  $D^-$  and is the key to the interpretation of results in the present paper.

An expression for the computation of  $R(DA^-)$  is obtained by addition of (7) to the equation,

$$\Delta G^m(A^-) = \Delta G_H + \Delta G(A^-) - R(A_*), \quad (9)$$

for photoelectron emission by aqueous solutions of the reduced form  $A^-$  of the acceptor A. The change of free energy  $\Delta G(A^-)$  pertains to the oxidation of  $A^-$  whereas  $\Delta G(A)$  is defined for reduction of A. Hence,  $\Delta G(A^-) = -\Delta G(A)$ . It will also be assumed that the quantities  $R(A_*^-)$  and  $R(A_*)$  pertaining to CT and emission, respectively, are equal. Conditions under which this assumption is justified are discussed in sec. 3. Thus,

$$\Delta G^m(A^-) - \Delta G^{CT} = 2\Delta G_H - \Delta G^m(D^-) + 2\Delta G(A^-) + R(DA^-). \quad (10)$$

The quantity  $R(DA^-)$  can be computed from (10), and the distance  $r_{DA}$  between the centers of the donor and acceptor is obtained from (3). This is significant because  $r_{DA}$  is obtained from experimental free energies without the introduction of the radii  $a_D$  and  $a_A$ . The assumption of outer-sphere contact in electron transfer in solution may be verified in this way.

## 2. Charge transfer from donor to solvent

Equation (7) will be tested for anions and cations as donors. The species  $A^-$  produced by CT is the hydrated electron, and  $\Delta G(A) = 2.77$  eV. One has [6]  $\Delta G^m(D^-) \approx E^t$ , where  $E^t$  is the experimental threshold energy for emission by an aqueous solution of the donor. Values of  $E^t$  were taken from [6-8] except for  $\text{Fe}(\text{CN})_6^{4-}$  ( $E^t = 6.6$  eV [9]). One also sets  $\Delta G^{CT}$  equal to the photon energy  $E^{\max}$  (from [10]) at the maximum of the CT absorption band of the donor.  $E^{\max}$  for  $\text{H}_2\text{PO}_4^-$  and  $\text{HPO}_4^{2-}$  were obtained by extrapolation and were not included in the least-square fitting discussed below. The photon energy for  $\text{SO}_3^{2-}$  was not used either since it corresponds to an extinction coefficient of  $10^3$  and not to the band maximum.

Least-square fitting yielded  $E^t - 4.50 = -0.03 + 1.016(E^{\max} - 2.77)$ , with  $E^t$  and  $E^{\max}$  in electronvolts. Hence,  $E^t - 4.50 \approx E^{\max} - 2.77$ . One expects from (7) that these two quantities differ by  $R$  provided that one sets  $\Delta G^m(D^-) = E^t$  and  $\Delta G^{CT} = E^{\max}$ , and the reorganization terms in (8) therefore nearly cancel out. Since one has  $R_{in}(A_\star^-) < 0$ , it follows that  $R(DA^-) > -R_{out}(A_\star^-)$  or, according to (3),  $r_{DA} < 2a_A$ . Taking  $r_{DA} = a_A + a_D$ , one must have  $a_D < a_A$  to account for the value  $R \approx 0$ . This is not surprising since model calculations [11] yield an equilibrium cavity radius of ca. 0.37 nm for the hydrated electron. This radius may be compared with the crystallographic radii ( $< a_D$ ) of ca. 0.2 nm for the donors of Fig. 1. The model leading to eq. (3) is approximate and so are radius values, but the preceding argument undoubtedly accounts for the near cancellation of terms in (8).

The quantities  $\Delta G_H$  and  $\Delta G(A)$  in (7) pertain to the final states of the electron, namely the vacuum level in emission and the level of the redox couple  $A/A^-$  in CT. Hence, emission and CT for the anions of Fig. 1 involve nearly the same free energies after correction for the difference in the final states of the

electron. This conclusion rests on the experimental values of  $E^t$  and  $E^{\max}$  without recourse to model calculations. Such model calculations for the CT process, although somewhat crude, in fact yield results in agreement with the present work, namely  $E^t \approx E^{\max} + 1.7$  (eV) [6].

The testing of eq. (7) for CT from cations to the solvent is more tentative than for anions because CT absorption bands of cations are hidden by other transitions. Only  $\text{Cr}^{2+}$  among the cations  $M^{2+}$  whose spectra are given in [12] exhibits a well-defined shoulder at ca. 4.6 eV [1]. Thus, one sets tentatively  $\Delta G_{CT} \approx 4.6$  eV for  $\text{Cr}^{2+}$ . The CT spectra of other cations can be characterized by the photon energy  $E^{\text{low}}$  [12] at which the extinction coefficient is 0.1. The plot (Fig. 2) with  $E^{\text{low}}$  instead of  $E^{\max} - 2.77$  (eV) appears quite linear. Least square fitting yielded  $E^t - 4.50 = -1.63 + 0.996 E^{\text{low}}$ , where the energies are in electronvolts. One has  $E^{\text{low}} = 3.35$  eV and  $\Delta G_{CT} \approx 4.6$  eV (see above) for  $\text{Cr}^{2+}$ . The intercept of the plot of  $E^t - 4.50$  (eV) against  $\Delta G_{CT}$  therefore is  $-1.63 - 0.996(4.6 - 3.35) = -2.88$  eV if one assumes that  $\Delta G_{CT} - E^{\text{low}}$  is the same for the cations of Fig. 2. If this is indeed the case, it may be concluded that the quantities  $E^t - 4.50$  and  $\Delta G_{CT} - 2.77$  (eV) are equal within a few tenths of electronvolt for each of the cations of Fig. 2. This conclusion is obviously tentative in view of the limited spectroscopic evidence.

### 3. Charge transfer from solvent to acceptor

Absorption spectra are available for CT from the solvent (water) to a few oxidizing species [13,14]. The photon energies  $E^{\max}$  of the CT absorption bands are listed in Table 1 together with  $\Delta G(A)$  [15,16]. Values of  $R$  in Table 1 were computed from (7) for  $\Delta G^m(D^-) = E^t = 10.06$  eV [7] for water.

The results for  $\text{Fe}^{3+}$  and  $\text{Ce}^{4+}$  will be interpreted first. The value  $R(DA^-) = 1.58$  eV was computed from (10) for  $\Delta G^m(A^-) = E^t = 7.35$  eV [5]



and  $\Delta G(A) = 0.77$  eV [16] for emission by  $\text{Fe}^{2+}$ . One deduces  $r_{\text{DA}} = 0.50$  nm from (3) for  $R(\text{DA}^-) = 1.58$  eV. This length is nearly equal to the distance for contact between a water molecule and a hexaquo ion  $\text{Fe}^{3+}$ , namely  $0.138 + (0.064 + 2 \times 0.138) = 0.48$  nm (0.064 and 0.138 nm, crystallographic radii of  $\text{Fe}^{3+}$  and  $\text{H}_2\text{O}$ , respectively). The agreement is remarkable, but experimental errors and the approximate character of eq. (3) and the relationship  $R(A_\star) = R(A_\star^-)$  should not be overlooked.

The value  $R = -0.50$  eV in Table 1 for  $\text{Fe}^{3+}$  is close to the inner-sphere reorganization free energy of  $-0.46$  eV for  $\text{Fe}^{3+}/\text{Fe}^{2+}$  electron transfer computed in [17] from the bond-stretching model. This agreement suggests (cf. eq. (8)) that  $R_{\text{out}}(A_\star^-) \approx -R(\text{DA}^-)$  within experimental and systematic errors of a few tenths of electronvolt. Likewise, one has  $R = -0.15$  eV for  $\text{Ce}^{4+}$  (Table 1) versus an inner-sphere term of  $-0.10$  eV in [17] for  $\text{Ce}^{4+}/\text{Ce}^{3+}$ . It is concluded (eq. (7)) that the energetics are nearly the same for emission by water and CT to  $\text{Fe}^{3+}$  or  $\text{Ce}^{4+}$  once  $E^t$  for emission and  $E^{\text{max}}$  for CT are corrected for reorganization and the difference of the final states of the electron.

The other acceptors in Table 1 will now be examined. Values of  $R(\text{DA}^-)$  between 0.6 and 0.9 eV are obtained for the halogens by application of (10), and the corresponding distance  $r_{\text{DA}} \approx 1$  nm computed (from (3)) is approximately twice too long (see above). It is concluded that the equality,  $R(A_\star^-) = R(A_\star)$ , introduced in the derivation of (10) does not hold for the halogens. Thus, the reorganization free energies for the processes  $A^- \rightarrow A_\star$  and  $A \rightarrow A_\star^-$  are not the same. This is to be expected from consideration of the curves R and P representing the total (electron involved in CT, ions, solvent) energy U against generalized reaction coordinate for the reactants and products of CT [2]. The energy  $U(R \rightarrow P)$  for the vertical transition from the

ground state of R to P is generally different from the energy  $U(P \rightarrow R)$  for the vertical transition from the ground state of P to R. This comparison must be made after correction for the change of free energy for CT. The difference between the two energies for vertical transition increases (in absolute value) with the difference between the coordinates of the ground states of R and P.

The radii  $a(\text{Fe}^{3+})$  and  $a(\text{Fe}^{2+})$  are nearly the same, and the near equality,  $R(\text{Fe}_*^{2+}) \approx R(\text{Fe}_*^{3+})$ , assumed in the calculation of  $R(\text{DA}^-)$  from (10) is consistent with the foregoing discussion of vertical transitions. Conversely, the atomic radii of the halogen atoms are definitely smaller than the corresponding crystallographic radii of the halide ions, and it is not surprising that  $|R(\text{A}_*^-)|$  for  $\text{A} \rightarrow \text{A}_*^-$  (CT) is different from  $|R(\text{A}_*)|$  for  $\text{A}^- \rightarrow \text{A}_*$  (emission). The high values of  $|R|$  for the halogens in Table 1 thus are accounted for by the higher (in absolute value) inner-sphere reorganization free energy for CT than for emission. The same conclusion holds for H and OH in Table 1. The increase of  $|R|$  from the inner-sphere term is particularly large for OH because vibrational excitation of both solvent and radical is involved.

#### 4. Donor-acceptor charge transfer in the 7 to 10 eV range

Equation (7) will be applied to the recently studied [8,18] CT processes between a wide variety of donors and acceptors in aqueous solution upon photon absorption in the 7 to 10 eV range. These results will be interpreted in terms of optical CT, but the analysis also holds if one assumes a scavenging mechanism. The confined volume model in [18], however, supports interpretation based on optical CT. The occurrence of CT was inferred from the lowering of the yield for photoelectron emission by the donor in presence of the acceptor. This method of detection of CT can be applied provided that the condition  $\Delta G^{\text{CT}} \leq \Delta G^{\text{m}}(\text{D}^-)$  holds, that is ((7) and (8)),

$$\Delta G_H - \Delta G(A) + R \geq 0. \quad (11)$$

If  $R = 0$ , condition (11) becomes  $\Delta G(A) \leq \Delta G_H$ , that is,  $\Delta G(A) \leq 4.5$  eV.

Strongly reducing species therefore may be produced by CT from the donor. CT was observed [18], for instance, with the following acceptors ( $\Delta G(A)$  in eV [16,19] between parentheses):  $H^+$  (2.1),  $Fe^{2+}$  (3.1),  $Ba^{2+}$  ( $4.85 \pm 0.25$ ).

CT to the alkali metal cations was not detected in [18] although the free energy change,  $\Delta G(A) \approx 3.5$  to  $3.6$  eV [19], is much more favorable to CT than for the CT-exhibiting  $Ba^{2+}$  ion ( $4.85 \pm 0.25$  eV). This seemingly paradoxical result can be understood by consideration of the term  $R$  in (11). Assume to simplify matters that  $R \approx R_{in}(A^+)$  since the other two terms contributing to  $R$  in (8) should nearly cancel out. Thus, one expects  $R_{in}(Na^+)$  to be significantly different from  $R_{in}(Ba^{2+})$  because CT to  $Na^+$  as acceptor would involve a much more pronounced change in size upon reduction to  $Na$  than CT to  $Ba^{2+}$  as acceptor. This difference in  $R$  more than compensates the difference in  $\Delta G(A)$ , and CT is only observed with  $Ba^{2+}$ . In contrast to the case of  $Na^+$ , CT to  $Tl^+$  is observed despite the marked increase in radius from ion to atom. CT is possible in that case because  $\Delta G(A)$  for the reduction of  $Tl^+$  is sufficiently low (1.9 eV) to offset the large inner-sphere term in  $R$ .

#### Acknowledgment

This work was supported by the Office of Naval Research and the National Science Foundation.

References

- [1] R. D. Cannon, Adv. Inorg. Chem. Radiochem. 21 (1978) 179.
- [2] R. D. Cannon, Electron transfer reactions (Butterworths, London, 1980) pp. 175-222, 267-297.
- [3] R. A. Marcus, J. Chem. Phys. 24 (1956) 966; 43 (1965) 679.
- [4] N. S. Hush, Trans. Faraday Soc. 57 (1961) 557.
- [5] P. Delahay, K. von Burg and A. Dziedzic, Chem. Phys. Letters 79 (1981) 157.
- [6] K. von Burg and P. Delahay, Chem. Phys. Letters 78 (1981) 287.
- [7] P. Delahay and K. von Burg, Chem. Phys. Letters 83 (1981) 250.
- [8] K. von Burg and P. Delahay, Chem. Phys. Letters, in press.
- [9] P. Delahay, unpublished results.
- [10] M. J. Blandamer and M. F. Fox, Chem. Rev. 70 (1970) 59, and references therein.
- [11] N. R. Kestner in Electron-solvent and anion-solvent intereactions, L. Kevan and B. C. Webster, eds. (Elsevier, Amsterdam, 1976) pp. 1-43.
- [12] F. S. Dainton and D. G. L. James, Trans. Faraday Soc. 54 (1958) 649.
- [13] E. Rabinowitch, Rev. Mod. Phys. 14 (1942) 112.
- [14] A. Treinin and E. Hayon, J. Am. Chem. Soc. 97 (1975) 1716, and references therein.
- [15] V. M. Berdnikov and N. M. Bazhin, Russ. J. Phys. Chem. 44 (1970) 395.
- [16] W. M. Latimer, Oxidation potentials (Prentice-Hall, New York, 1952).
- [17] J. M. Hale in Reactions of molecules at electrodes, N. S. Hush, ed. (Wiley-Interscience, New York, 1971) pp. 229-257.
- [18] K. von Burg and P. Delahay, Chem. Phys. Letters 83 (1981) 199.
- [19] J. H. Baxendale and R. S. Dixon, Z. phys. Chem. N.F. 43 (1964) 161.

Table 1

Charge transfer from water to acceptor

Acceptor	$E^{\max}$ a) (eV)	$-\Delta G(A)$ b) (eV)	$-R$ (eV)	$-R_{in}(A_*)$ c) (eV)
H	>6.2	-0.1	>0.5	-
OH	5.40	1.9	1.74	-
Cl	3.95	2.55	0.94	-
Br	4.50	2.0	0.94	-
I	4.85	1.4	0.69	-
Fe <sup>3+</sup>	5.29	0.77	0.50	0.46
Ce <sup>4+</sup>	4.00	1.70	0.14	0.10

a) From [13,14].  $E^{\max}$  for H higher than upper limit (6.2 eV) of explored range of photon energies ( $E^{\text{low}} \approx 5.0$  eV).

b) From [15,16].

c) From [17].

## Captions to Figures

Fig. 1. Plot (eq. (7)) for CT from anion to solvent (water).  $E^t$ , threshold energy for photoelectron emission by anions;  $E^{\max}$ , photon energy at maximum of CT-absorption band of the anions.

Fig. 2. Plot for CT from cation to solvent (water).  $E^{\text{low}}$ , photon energy at which the extinction coefficient is 0.1 on the cation absorption spectra from [12].




FIG. 1




FIG. 2



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Western Regional Office Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Eastern Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
		Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.  
Copies

Mr. James Kelley  
DTNSRDC Code 2803  
Annapolis, Maryland 21402

1

Mr. A. M. Anzalone  
Administrative Librarian  
PLASTEC/ARRADCOM  
Bldg 3401  
Dover, New Jersey 07801

1

TECHNICAL REPORT DISTRIBUTION LIST, 0515

	<u>No. Copies</u>		<u>No. Copies</u>
Professor R. Wilson Department of Chemistry, 8-014 University of California, San Diego La Jolla, California 92093	1	Dr. B. Vonnegut State University of New York Earth Sciences Building 1400 Washington Avenue Albany, New York 12203	1
Professor C. A. Angell Department of Chemistry Purdue University West Lafayette, Indiana 47907	1	Dr. Hank Loos Laguna Research Laboratory 21421 Stans Lane Laguna Beach, California 92651	1
Professor P. Meijer Department of Physics Catholic University of America Washington, D.C. 20064	1	Dr. John Latham University of Manchester Institute of Science & Technology P.O. Box 88 Manchester, England M601QD	1
Dr. S. Greer Chemistry Department University of Maryland College Park, Maryland 20742	1		
<del>Professor P. Delahay</del> <del>New York University</del> <del>100 Washington Square East</del> <del>New York, New York 10003</del>	1		
Dr. T. Ashworth Department of Physics South Dakota School of Mines & Technology Rapid City, South Dakota 57701	1		
Dr. G. Gross New Mexico Institute of Mining & Technology Socorro, New Mexico 87801	1		
Dr. J. Kassner Space Science Research Center University of Missouri - Rolla Rolla, Missouri 65401	1		
Dr. J. Telford University of Nevada System Desert Research Institute Lab of Atmospheric Physics Reno, Nevada 89507	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Paul Delahay Department of Chemistry New York University New York, New York 10003	1	Dr. P. J. Hendra Department of Chemistry University of Southampton Southampton SO9 5NH United Kingdom	1
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106	1	Dr. Sam Perone Department of Chemistry Purdue University West Lafayette, Indiana 47907	1
Dr. D. N. Bennion Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
Dr. J. J. Auburn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research & Technology Department Naval Surface Weapons Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. T. Katan Lockheed Missiles & Space Co, Inc. P.O. Box 504 Sunnyvale, California 94088	1	Dr. J. Boechler Electrochimica Corporation Attention: Technical Library 2485 Charleston Road Mountain View, California 94040	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063	1
Dr. B. Brummer EIC Incorporated 55 Chapel Street Newton, Massachusetts 02158	1	Dr. H. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1	Dr. R. P. Van Duyne Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. B. Stanley Pons Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6G 2G2	1
Larry E. Plew Naval Weapons Support Center Code 30736, Building 2906 Crane, Indiana 47522	1	Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824	1
S. Rubv DOE (STOR) 600 E Street Washington, D.C. 20545	1	Dr. R. David Rauh EIC Corporation 55 Chapel Street Newton, Massachusetts 02158	1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	1	Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1	Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton 509 5NH England	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1	Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California	1	Dr. R. A. Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910	1	Mr. James R. Moden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. R. Nowak Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Dr. Bernard Spielvogel U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709	1
Dr. John F. Houlihan Shenango Valley Campus Pennsylvania State University Sharon, Pennsylvania 16146	1	Dr. Denton Elliott Air Force Office of Scientific Research Bolling AFB Washington, DC 20332	1
Dr. D. F. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201	1	Dr. David Aikens Chemistry Department Rensselaer Polytechnic Institute Troy, NY 12181	1
Dr. D. H. Whitmore Department of Materials Science Northwestern University Evanston, Illinois 60201	1	Dr. A. P. B. Lever Chemistry Department York University Downsview, Ontario M3J1P3 Canada	1
Dr. Alan Bewick Department of Chemistry The University Southampton, SO9 5NH England	1	Mr. Maurice F. Murphy Naval Sea Systems Command 63R32 2221 Jefferson Davis Highway Arlington, VA 20360	1
Dr. A. Himy NAVSEA-5433 NC #4 2541 Jefferson Davis Highway Arlington, Virginia 20362	1	Dr. Stanislaw Szpak Naval Ocean Systems Center Code 6343 San Diego, CA 95152	1
Dr. John Kincaid Department of the Navy Strategic Systems Project Office Room 901 Washington, DC 20376	1	Dr. Gregory Farrington Department of Materials Science & Engineering University of Pennsylvania Philadelphia, PA 19104	1
M. L. Robertson Manager, Electrochemical Power Sonics Division Naval Weapons Support Center Crane, Indiana 47522	1	Dr. Bruce Dunn Department of Engineering & Applied Science University of California Los Angeles, CA 90024	1
Dr. Elton Cairns Energy & Environment Division Lawrence Berkeley Laboratory University of California Berkeley, California 94720	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

No.  
Copies

Dr. Micha Tomkiewicz  
Department of Physics  
Brooklyn College  
Brooklyn, NY 11210 1

Dr. Lesser Blum  
Department of Physics  
University of Puerto Rico  
Rio Piedras, PR 00931 1

Dr. Joseph Gordon II  
IBM Corporation  
K33/281  
5600 Cottle Road  
San Jose, CA 95193 1

Dr. Robert Somoano  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, CA 91103 1